

R package to handle  
Archimax or any user-defined continuous  
copula construction:  
*acopula*

Tomáš Bacigál

Slovak University of Technology in Bratislava  
Slovakia

AGOP 2013


# Outline

- alternatives
- Archimax and generic copulae definition lists
- probability functions
- estimation
- testing
- utilities
- conclusion and further plans


# Alternatives

- Mathematica 8
- Matlab
- Excel
  - Hoadley Finance Add-in
  - Vose Model Risc
- S-Plus
  - S+ Finmetrics/EVANESCE
- XploRe, ...
- R
  - copula (nacopula)
  - CDVine/VineCopula
  - CopBasic, fCopulae, copulaedas, HAC, fgac, sbgcop, pencopula, penDvine, vines, ...
  - acopula

# acopula – building blocks



# acopula – building blocks



# acopula – building blocks

## generator

- Ali-Mikhail-Haq
- Clayton
- Frank
- Gumbel-Hougaard
- Joe
- log

## dependence function

- Galambos
- Gumbel-Hougaard
- Husler-Reiss
- Tawn
- 1, max
- (general) convex combination

## copula

- Farlie-Gumbel-Morgenstern
- Gumbel-Hougaard
- Normal
- Plackett
- product

```
> pCopula(data=c(0.2, 0.3), generator=genGumbel(), gpars=3.5)
```

```
> pCopula(data=c(0.2, 0.3), copula=copGumbel(), pars=3.5)
```

```
> pCopula(data=c(0.2, 0.3), gen=genLog(), depfun=depGumbel(), dpars=3.5)
```

```
[1] 0.1723903
```

# Probability functions

```
> gG <- genGumbel(parameters=3.5)
```

- $P(X < x \ \& \ Y < y) = ?$

```
> pCopula(c(x, y), gG)
```

- $P(X < ? \ \& \ Y < y) = p$

```
> pCopula(c(p, y), gG, qua=1)
```

```
> pCopula(c(NA, y), gG, qua=1,
```

```
+ prob=p)
```

```
> qCopula(y, qua=1, prob=p, gG)
```

- $P(X < x \ | \ Y = y) = ?$

```
> cCopula(c(x, y), con=2, gG)
```

- $P(X < ? \ | \ Y = y) = p$

```
> cCopula(c(p, y), con=2, gG, qua=1)
```

```
> pCopula(c(NA, y), con=2, gG, qua=1,
```

```
+ prob=p)
```

```
> qCopula(y, qua=1, prob=p, con=2,
```


```
+ gG)
```

# Probability functions

```
> gG <- genGumbel(parameters=3.5)
```


- density

```
> dCopula(c(x, y), gG)
```



- random sampling


```
> rCopula(100, dim=2, gG)
```



- empirical copula CDF

```
> pCopulaEmpirical(c(x, y), base=sample)
```

# Estimation



```
> eCopula(sample, gen=genClayton(), dep=depGumbel(),  
+ technique="ML", procedure="optim", method="L-BFGS-B")
```


```
generator parameters: 0.09357958
```

```
depfun parameters: 3.52958
```

```
ML function value: 82.63223
```

```
convergence code: 0
```

# Estimation



Note: Not all copula families has closed-form relation between its parameters and a correlation coefficient, yet it can be approximated by, e.g., 4-parametric Parreto CDF which is easily invertible  $\rightarrow$  max. approx. error at worst  $\pm 0.01$  mostly  $\pm 0.001$

# Estimation

- Kendall's tau

$$\tau = 4 \iint_{[0,1] \times [0,1]} C(u,v)c(u,v) \, dudv - 1$$

- Spearman's rho

$$\rho = 12 \iint_{[0,1] \times [0,1]} C(u,v) \, dudv - 3$$

# Estimation

- Kendall's tau

$$\tau = 4 \iint_{[0,1] \times [0,1]} C(u,v)c(u,v) \, du dv - 1$$

- Gumbel  $\tau = (\theta - 1) / \theta$

- Clayton  $\tau = \theta / (\theta + 2)$

- FGM  $\tau = 2\theta / 9$

Not many have a closed form, but the relation can be approximated.

# Estimation

- Kendall's tau

$$\tau = 4 \iint_{[0,1] \times [0,1]} C(u,v)c(u,v) \, du \, dv - 1$$

- Gumbel  $\tau = (\theta - 1) / \theta$


- Clayton  $\tau = \theta / (\theta + 2)$

- FGM  $\tau = 2\theta / 9$

Not many have a closed form, but the relation can be approximated.

For example

- Frank



# Estimation

- Kendall's tau

$$\tau = 4 \iint_{[0,1] \times [0,1]} C(u,v)c(u,v) \, du \, dv - 1$$

- Gumbel  $\tau = (\theta - 1) / \theta$

- Clayton  $\tau = \theta / (\theta + 2)$

- FGM  $\tau = 2 \theta / 9$

Not many have a closed form, but the relation can be approximated.

For example


- Frank  $\tau = \text{CDF}(\theta)$

$$k = 6.1, \alpha = 1.1, \gamma = 0.8,$$

$$\mu = 0$$

## Pareto type IV distribution

$$CDF(x) = \begin{cases} 1 - \left(1 + \left(\frac{x - \mu}{k}\right)^{\frac{1}{\gamma}}\right)^{-\alpha} & x \geq \mu \\ 0 & \text{otherwise} \end{cases}$$



# Estimation

- Kendall's tau

$$\tau = 4 \iint_{[0,1] \times [0,1]} C(u,v)c(u,v) \, du \, dv - 1$$

- Gumbel  $\tau = (\theta - 1) / \theta$

- Clayton  $\tau = \theta / (\theta + 2)$

- FGM  $\tau = 2 \theta / 9$

Not many have a closed form, but the relation can be approximated.

For example


- Frank  $\tau = \text{CDF}(\theta)$

$$k = 6.1, \alpha = 1.1, \gamma = 0.8,$$

$$\mu = 0$$

## Pareto type IV distribution

$$CDF(x) = \begin{cases} 1 - \left(1 + \left(\frac{x - \mu}{k}\right)^{\frac{1}{\gamma}}\right)^{-\alpha} & x \geq \mu \\ 0 & \text{otherwise} \end{cases}$$



# Testing

- goodness-of-fit

```
> gCopula (sample, cop=copNormal(),  
+ etechnique="ML", eprocedure="optim", ncores=1, N=100)  
|=====| 100%
```

Blanket GOF test based on Kendall's transform

statistic	q95	p.value
0.1195500	0.1658125	0.1800000

-----

data: sample

copula: normal

estimates:

pars	fvalue
------	--------

0.9155766	80.3420886
-----------	------------

# Testing

- goodness-of-fit
- equality of 2 copulas

```
> sampleC1 <- rCopula(n=100, dim=2, gen=genClayton(), gpars=5)
> gCopula(list(sample, sampleC1), ncores=1, N=100)

|=====| 100%
```

Test of equality between 2 empirical copulas

statistic	q95	p.value
0.09791672	0.52893392	0.66000000

-----

data: sample sampleC1

copula:

estimates:

NULL

# Testing

- goodness-of-fit
- equality of 2 copulas
- validity of copula properties
  - monotonicity, annihilator and neutral element

```
> isCopula(gen=genGumbel(lower=0), dim=3, glimits=list(0.5, 2),  
+ dagrid=10, pgrid=4, tolerance=1e-15)
```

```
Does the object appears to be a copula(?): FALSE
```

```
Showing 2 of 2 issues:
```

	dim	property	value	gpar
1	2	monot	-0.1534827	0.5
2	3	monot	-0.1402209	0.5

# Utilities

- numerical derivative

```
> fun <- function(x,y,z) x^2*y*exp(z)
> nderive(fun,point=c(0.2,1.3,0),order=c(2,0,1),
+ difference=1e-04,area=0)
[1] 2.600004
```

# Utilities

- numerical derivative
- numerical integration

```
> nintegrate(function(x,y) mvtnorm::dmvnorm(c(x,y)),  
+ lower=c(-5.,-5.), upper=c(0.5,1), subdivisions=30)
```

```
[1] 0.5807843
```

```
> #Better alternative:
```

```
> cubature::adaptIntegrate(mvtnorm::dmvnorm,  
+ lowerLimit=c(-5.,-5.), upperLimit=c(0.5,1))$integral
```

```
[1] 0.5817578
```

```
> pnorm(0.5)*pnorm(1) #exact solution due to independence
```

```
[1] 0.5817583
```

# Conclusion

- arbitrary dimension (with `nderive` for  $\text{dim} > 2$ )
- conditional probability and quantile function
- generalization of Archimedean and EV class
- construction method of Pickand's dependence function
- test of equality between two empirical copulas
- numerical check of copula properties
- parallelized goodness-of-fit test based on Kendall's transform
- estimation based on inversion of correlation coefficients available for every 1-parameter copula family

# Further plans

- non-parametric estimation for multi-parameter families
- new construction methods
- more GoF test methods
- more copulas definition lists
- connection to other copula packages
- procedures for practical analysis
- bug fixing

Thank you for attention and new ideas.